



501162
TCT 195 03 / 00106

GB03 / 156



INVESTOR IN PEOPLE

The Patent Office
Concept House
Cardiff Road
Newport
South Wales
NP10 8QQ


REC'D	07 MAR 2003
WIPO	PCT

I, the undersigned, being an officer duly authorised in accordance with Section 74(1) and (4) of the Deregulation & Contracting Out Act 1994, to sign and issue certificates on behalf of the Comptroller-General, hereby certify that annexed hereto is a true copy of the documents as originally filed in connection with the patent application identified therein.

In accordance with the Patents (Companies Re-registration) Rules 1982, if a company named in this certificate and any accompanying documents has re-registered under the Companies Act 1980 with the same name as that with which it was registered immediately before re-registration save for the substitution as, or inclusion as, the last part of the name of the words "public limited company" or their equivalents in Welsh, references to the name of the company in this certificate and any accompanying documents shall be treated as references to the name with which it is so re-registered.

In accordance with the rules, the words "public limited company" may be replaced by p.l.c., plc, P.L.C. or PLC.

Re-registration under the Companies Act does not constitute a new legal entity but merely subjects the company to certain additional company law rules.



Andrew Gensley

Signed

Dated 7 October 2005


**PRIORITY
DOCUMENT**

SUBMITTED OR TRANSMITTED IN
COMPLIANCE WITH RULE 17.1(a) OR (b)

BEST AVAILABLE COPY

Patents Form 1/77

Patents Act 1977
(Section 16)



1/77

The Patent Office
Cardiff Road
Newport
Gwent NP9 1RH

Request for grant of a patent

(See the notes on the back of this form. You can also get an explanatory leaflet from the Patent Office to help you fill in this form)

1. Your reference 27.17.77359/00122NOV02 E765434-2 D00027
P01/7700 0.00-0227239.1

2. Patent application number 0227239.1
(The Patent Office will fill in this part)

3. Full name, address and postcode of the
or of each applicant (underline all surnames)
Dynal Biotech ASA
Postboks 114 Smestad
N-0309 Oslo
Norway

Patents ADP number (if you know it)

If the applicant is a corporate body, give
country/state of incorporation

4. Title of the invention DNA/RNA and protein isolation from a
single sample

5. Name of your agent (if you have one) Frank B. Dehn & Co.

"Address for service" in the United Kingdom
to which all correspondence should be sent
(including the postcode)

179 Queen Victoria Street
London
EC4V 4EL

Patents ADP number (if you know it) 166001

6. If you are declaring priority from one or more earlier patent applications, give the country and the date of filing of the or of each of these earlier applications and (if you know it) the or each application number	Country	Priority application number (if you know it)	Date of filing (day / month / year)

7. If this application is divided or otherwise derived from an earlier UK application, give the number and the filing date of the earlier application	Number of earlier application	Date of filing (day / month / year)

8. Is a statement of inventorship and of right
to grant of a patent required in support of
this request? (Answer 'Yes' if:
a) any applicant named in part 3 is not an inventor, or
b) there is an inventor who is not named as an
applicant, or
c) any named applicant is a corporate body.
See note (d))

yes

08/6/65500

Patents Form 1/77

9 Enter the number of sheets for any of the following items you are filing with this form. Do not count copies of the same document

Continuation sheets of this form 0

Description 47

Claim(s) 0

Abstract 0

Drawing(s) 12 *212*

10. If you are also filing any of the following, state how many against each item.

Priority documents 0

Translations of priority documents 0

Statement of inventorship and right to grant of a patent (Patents Form 7/77) 0

Request for preliminary examination and search (Patents Form 9/77) 0

Request for substantive examination (Patents Form 10/77) 0

Any other documents (please specify) 0

11. I/We request the grant of a patent on the basis of this application.
Frank B. Dehn & Co.
Signature Date 21 November 2002

12. Name and daytime telephone number of person to contact in the United Kingdom
Hanna Dzieglewska
020 7206 0600

Warning

After an application for a patent has been filed, the Comptroller of the Patent Office will consider whether publication or communication of the invention should be prohibited or restricted under Section 22 of the Patents Act 1977. You will be informed if it is necessary to prohibit or restrict your invention in this way. Furthermore, if you live in the United Kingdom, Section 23 of the Patents Act 1977 stops you from applying for a patent abroad without first getting written permission from the Patent Office unless an application has been filed at least 6 weeks beforehand in the United Kingdom for a patent for the same invention and either no direction prohibiting publication or communication has been given, or any such direction has been revoked.

Notes

- If you need help to fill in this form or you have any questions, please contact the Patent Office on 0645 500505.
- Write your answers in capital letters using black ink or you may type them.
- If there is not enough space for all the relevant details on any part of this form, please continue on a separate sheet of paper and write "see continuation sheet" in the relevant part(s) of the form. Any continuation sheet should be attached to this form.
- If you have answered 'Yes', Patents Form 7/77 will need to be filed.
- Once you have filled in the form you must remember to sign and date it.
- For details of the fee and ways to pay please contact the Patent Office.

77359/001.619

Method

5 The present invention relates to a method of
analysing a single sample for DNA, mRNA and protein
expression. This is advantageous in correlating
information relating to both mRNA and protein expression
to genomic information.

10 The human genome is now more or less mapped and the
gene number has been found to be much lower than first
estimated. This shows that human complexity does not
lie in the number of genes, but in how the genes are
utilized and gene products combined and modified. Thus,
the focus on studying the gene products has become more
15 significant.

It is known that only a fraction of the genes are
transcribed in a single cell or cell type, and that one
gene can give rise to many different proteins. This is
partly due to alternative reading frames, alternative
20 translational start/stop codons and alternative splicing
of mRNA. It is also known that the relative
quantitative levels of mRNA and protein do not always-
or seldom correspond. One mRNA species can also give
rise to different proteins because of post-translational
25 modifications.

Thus an individual's genome (the collection of all
genes), can be regarded as representing the genetic
"possibilities" of an individual, while the
transcriptome (the collection of mRNAs transcribed from
30 the genome) represents what might possibly happen. The
proteome, however (i.e. the total collection of all
proteins translated from the transcriptome and modified
post-translationally) represents reality, i.e. what
really happened.

35 Accordingly, all three levels, DNA, RNA and
protein, give information which is valuable for
different reasons. The DNA or genotype gives important

information about genetic pre-dispositions and acquired mutations/local rearrangements. Both mRNA and protein profiles generate "molecular portraits" of a state/stage or disease, and may also be used for staging and
5 monitoring of the disease development and treatment. As opposed to the DNA, both mRNA and protein profiles represent "snap shots" of the cell's biology, since they are continuously changing in response to the surrounding environment. Due to regulatory mechanisms acting both
10 at the transcriptional, translational and post-translational levels, mRNA and protein levels do not always correlate. It is therefore crucial to study both mRNA and protein from the same sample.

Thus, as mentioned above, there is however not
15 necessarily a 1:1 correlation between mRNA and protein levels. If protein levels and corresponding mRNA levels are compared, then for every protein for which the ratio of mRNA and protein is not 1:1 then this protein is subject to some form of interesting post transcriptional
20 regulation. mRNA/protein ratios for specific genes are often shifted during disease conditions. To be able to study regulatory mechanisms and to unravel the reasons behind such a shift in mRNA/protein ratios, it is crucial to isolate mRNA and protein from the same
25 sample. A novel and advantageous method for carrying out such isolation is presented herein.

Variation in protein or mRNA levels between individuals, for example in an individual with a disease state compared with a "normal" patient, are often due to
30 single nucleotide polymorphisms (SNPs) or other forms of mutation or genetic predisposition in their genes (i.e. their DNA) or their regulatory regions. For this reason it is advantageous to be able to isolate DNA and protein and preferably DNA, RNA and protein from the same sample
35 to enable direct comparison. Such methods are also provided by the present invention.

In particular, it has now been found that nucleic

acid and protein, and in particular DNA, RNA and protein can be isolated from a single sample in a form suitable for downstream manipulation and analysis, by a simple and easy to perform procedure. This isolation method

5 involves binding nucleic acid and protein or preferably DNA, RNA and protein from the same sample on to distinct solid supports or areas of solid supports so that the isolated components can be analysed separately.

Advantageously these methods can be combined with a

10 further initial step involving the specific isolation of one or more particular cell types from the sample before the isolation of nucleic acid and protein is carried out. In particular, it has been shown that DNA (e.g. genomic DNA), RNA (e.g. mRNA) and proteins may be

15 isolated from a single sample with a high, documented purity.

In one aspect, the present invention thus provides a method of isolating nucleic acid and protein from the same sample, said method comprising contacting said

20 sample with solid supports, whereby the nucleic acid and protein components in said sample become bound to distinct solid supports.

The nucleic acid to be isolated may be DNA, RNA or any naturally occurring modification thereof, and

25 combinations thereof. Thus, the entire nucleic acid based component of a sample can be isolated in the method of the invention. In a preferred embodiment of the invention however DNA and/or RNA are separately isolated from the same sample onto distinct solid

30 supports.

Thus, in preferred embodiments the present invention provides a method of isolating RNA and protein or DNA and protein from the same sample. In especially preferred embodiments the invention provides a method of

35 isolating DNA, RNA and protein from the same sample.

In embodiments and steps of the methods of the invention, where the nucleic acid to be isolated from

the sample is DNA, preferably the DNA is cDNA or more preferably genomic DNA and may be in a single or double stranded or in any other form. In certain embodiments a specific DNA molecule, e.g comprising a gene encoding a specific protein may be isolated.

In embodiments and steps of the methods of the invention, where the nucleic acid to be isolated from the sample is RNA, total RNA may be isolated or a particular form or subset of RNA may be isolated. In preferred embodiments of the invention mRNA is isolated from the sample. In certain embodiments a specific RNA molecule, e.g. a specific mRNA encoding a specific protein may be isolated.

In the methods of the present invention described herein the steps of isolating nucleic acid and protein onto distinct solid phases involving contacting the sample with said solid phases can be carried out sequentially or simultaneously. Similarly, in embodiments where it is desired to isolate protein, DNA and RNA as separate components then these isolations can be carried out in the same step or in separate sequential steps.

Preferably the required isolations are carried out in separate sequential steps which can, in turn, be carried out in any order. Thus, although in the preferred embodiments of the invention DNA is isolated in a first step, followed by RNA in a second step and protein in a third step the different components can be isolated in any order.

The term "distinct solid phases" as used herein includes the use of different solid supports (i.e. supports with different binding and surface properties) in different steps of the method of the invention (for example in embodiments where sequential steps are used to bind the desired nucleic acid components and proteins), different supports added in same step (for example in embodiments where the isolation of the

desired nucleic acid components and protein are carried out in the same step) and the use of supports with the same surface properties added in different steps, where binding of a particular nucleic acid or protein component is enabled by virtue of different incubation conditions. This term also includes different areas of a single solid support, which may for example be designed to have different surface properties such that particular nucleic acid and protein components bind to different areas of the same support.

As mentioned above, it is an important aspect of the invention that nucleic acid and protein are analysed from the same sample.

"Same sample" as used herein refers to the isolation of the appropriate nucleic acid and protein components from a single, i.e. undivided sample. In this way the method of the invention is distinguished from prior art methods which may involve obtaining a sample and dividing it into aliquots on which a separate analysis of DNA, RNA and protein (one for each aliquot) is carried out and the results compared. The fact that the multiple analysis of DNA, RNA and protein can be carried out on a single, undivided sample using the methods of the present invention is clearly advantageous and will allow for a more direct and accurate comparison between the various nucleic acid and protein components of the sample.

The samples suitable for use in the methods of the present invention may be any material containing nucleic acid and protein, including for example foods and allied products, clinical and environmental samples. However, the sample will generally be a biological sample, which may contain any viral or cellular material, including all prokaryotic or eukaryotic cells, viruses, bacteriophages, mycoplasmas, protoplasts and organelles. Such biological material may thus comprise all types of mammalian and non-mammalian animal cells, plant cells,

algae including blue-green algae, fungi, bacteria, protozoa etc. Representative samples thus include whole blood and blood-derived products such as plasma, serum and buffy coat, bone marrow and samples obtained from
5 other haematopoietic tissues, urine, faeces, cerebrospinal fluid or any other body fluids, tissues, cell cultures, cell suspensions etc. Preferably the sample is a live (e.g. a non-fixed sample), i.e. comprises viable cells, or at least cells which have not
10 been treated in any way.

Suitable samples may also include relatively pure, e.g. non-particulate/non-cell containing, starting materials which contain relatively pure nucleic acid and protein preparations, such as a PCR product, proteins
15 synthesised in vitro, or semi-pure preparations obtained by other nucleic acid and protein recovery processes.

The sample may be freshly obtained, or stored or treated in any desired or convenient way, for example by dilution, or adding buffer, or other solutions or
20 solvents, enzyme-containing solutions etc.), as long as the integrity of the nucleic acids and proteins, or the cells or other entities containing such nucleic acids and proteins within the original sample are maintained, i.e. are not substantially degraded.

25 It is further advantageous that a crude sample can be taken directly from its source, e.g. directly from a patient or from the environment (e.g. a direct clinical or environmental sample) and its protein and nucleic acid analysed using the methods of the invention.

30 The nucleic acid- and protein-containing sample may, generally speaking, simply be contacted with the appropriate solid supports under conditions whereby the nucleic acids or proteins as appropriate will become bound to the solid support. If necessary, e.g. if the
35 nucleic acid and protein to be isolated is not available for binding within the initial sample e.g. is contained within a biological particle as described above e.g. a

viral coat or a cell membrane or wall, the initial binding step may be preceded by one or more separate steps to free the nucleic acid and protein components, e.g. by disrupting structural components such as cell walls or to achieve lysis. Procedures for achieving this are well known in the art. Thus, for example, although some cells e.g. blood cells, may be lysed by reagents such as detergent alone, other cells, e.g. plant or fungal cells or solid animal tissues may require more vigorous treatment such as, for example, grinding in liquid nitrogen, heating in the presence of detergent, alkaline lysis in the presence of detergent. The main requirement is that procedures for freeing the nucleic acid and protein are chosen such that the particular nucleic acid and protein species which are to be isolated in the methods of the invention remain sufficiently intact, e.g. are not substantially degraded.

Preferably samples are 10 μ l to 100 ml in size, preferably from 200 μ l to 10 ml. The method of the invention may be used for small samples, e.g. less than 1 ml or for larger samples e.g. at least 2 ml, e.g. more than 5 ml or 10 ml or 50 ml. A major advantage of the present invention is that small sample volumes may be used, and in particular that protein may be isolated from such small sample volumes following, or concurrently with, DNA and/or RNA isolation. This feature makes the method of the invention particularly suitable for automation. Advantageously the sample volume as subjected to the nucleic acid/protein isolation procedures is 1 ml or less, e.g. 10 to 800 μ l, e.g. 20 to 500 μ l, or 50 to 200 μ l. For example, the method may be performed on a sample containing 1×10^4 to 1×10^6 cells, preferably $1-10 \times 10^5$ cells. The method described is highly scalable and can be used to detect DNA/RNA/protein in 1, 5, 10, 20, 200 or 2000 or more cells.

As mentioned above an optional and preferred

initial step once the sample has been obtained is to enrich the sample for particular populations, e.g. particular populations of cells or organelles for which the nucleic acid and protein is to be analysed.

5 Methods of enrichment are well known and documented in the art. Preferred methods involve the use of a solid phase, i.e. the binding of the population(s) in question to a solid phase and exemplary methods will be discussed in more detail below. In such methods, one or more
10 steps of positive and/or negative solid phase selection can be used, as appropriate, depending on the desired purity of sub-population required. Such techniques are well known and described in the art. Briefly, in a negative selection step, the cells/populations which
15 bind to the solid phases are removed from the sample to be analysed, whereas in a positive selection step the cells/populations which bind to the solid phases are retained in the sample to be analysed.

 Exemplary and preferred antibodies for use in the
20 isolation of specific cell populations are CD15 and CD45 antibodies (specific for leukocytes), CD14 antibodies (specific for monocytes), CD2, CD3, CD4 and CD8 antibodies (specific for T-cells), CD19 antibodies (specific for B-cells) and Anti Ber EP4 antibodies
25 (specific for epithelial cells and in particular circulating carcinoma cells). Other antigens may also be used as the basis for detection by binding to specific antibodies (or antibody fragments, derivatives etc.), as indeed may any other cell surface molecule
30 which may bind specifically to a binding partner or ligand.

 Other methods of isolating a desired cell population may also be used, as desired, for example Laser Capture Microdissection.

35 Where the method of the invention involves such a preliminary cell/organelle etc. isolation step, it will be appreciated that larger sample volumes may be used,

e.g. 1-50 ml, e.g. 2 to 20 ml or 5-10 ml, for example of a blood or other clinical sample.

Sample preparation is one of the bottlenecks in proteomics in that due to the variation in abundance and lack of specific amplification technology for proteins, it is advantageous to reduce sample complexity in order to increase resolution. Thus the optional step of the present invention by which a particular sample can be treated to specifically enrich for a particular specific population or populations of cells, organelles, etc., and thereby proteins is extremely advantageous. Thus, it can be seen that the methods of the invention allow the enrichment of a particular population of cells and the analysis of DNA, RNA and protein therefrom using the same single sample. The fact that this sample can be taken directly from a patient is a further advantage.

The feature of specifically or selectively isolating a desired population may be applied advantageously to comparing multiple (e.g. 2 or more, e.g. 2 to 10 or 2 to 6) populations from the same sample. Thus, for example, by the use of different appropriate binding partners specific for particular desired populations and coupled to different solid supports, different populations may be isolated, or separated from the same sample.

Thus, a preferred embodiment of the invention provides a method of isolating DNA, RNA, and protein from the same sample, said method comprising the following steps:

- a) isolating from said sample the populations from which the nucleic acid and protein is to be isolated and preferably isolating one or more specific populations;
- b) lysis of the populations, e.g. cell populations, to obtain a sample containing nucleic acid and protein in a free form available for binding to a solid support;

c) contacting sample with an appropriate solid phase under conditions whereby DNA, preferably genomic DNA, becomes bound to the solid support;

5 d) separation of this solid phase from the remainder of the sample;

e) contacting said remainder of the sample with an appropriate solid phase under conditions whereby RNA, preferably mRNA, becomes bound to the solid support;

10 f) separation of this solid phase from the remainder of the sample;

g) contacting said remainder of the sample with an appropriate solid phase under conditions whereby a specific protein or population of proteins becomes bound to the solid support;

15 h) separation of this solid phase from the remainder of the sample.

The method outlined above is a preferred embodiment of the invention and clearly these specific steps can be altered depending on the overall analysis it is wished to achieve to form further embodiments of the invention. In particular, some of the above discussed steps are optional and need not be included. For example, steps a) and b) are optional and whether or not they are carried out depends on the nature of the sample which is initially obtained. If this sample does not contain the nucleic acid and protein which it is desired to analyse in a free and released form, for example if the nucleic acid and protein is contained within cells or tissues or other biological packages, then generally steps a) and b) will be carried out.

20
25
30

In addition, the above described methods describes the sequential isolation of DNA, RNA and protein in separate steps. The isolation of the whole nucleic acid component as one fraction and protein, and the isolation of DNA and protein, and RNA and protein, are also within the scope of the invention and the steps of the method can be adapted as appropriate to fulfil this purpose.

35

For example, if it is desired to isolate the total nucleic acid and protein, step c) can be replaced by a step which involves contacting the sample with an appropriate solid phase under conditions whereby the total nucleic acid component, e.g. DNA and RNA, becomes bound to the solid support. Step e) would then be omitted. If it is desired to isolate DNA and protein then the RNA steps e) and f) can be omitted. Vice versa, if it is desired to isolate RNA and protein then the DNA steps c) and d) can be omitted.

In addition, the nucleic acid and protein isolation steps of the above described methods can be carried out in any order. Thus, although in a preferred embodiment the order is DNA followed by RNA followed by protein (if all three components are to be analysed), the steps may be carried in any order, e.g. the RNA or the protein may be isolated first. In methods where nucleic acid and protein are to be isolated, or DNA and protein, or RNA and protein are to be isolated, although in preferred embodiments the nucleic acid component is isolated first, the protein component may be isolated first.

In addition, in a further embodiment of the invention the desired nucleic acid and protein components can be isolated simultaneously from the same sample in the same step. This may be achieved in any appropriate manner but will generally be carried out by contacting the sample simultaneously with distinct solid supports which have different surface properties and are able to bind specifically to DNA (e.g. genomic DNA), RNA (e.g. mRNA), all nucleic acid species, or protein, as appropriate. Appropriate solid supports which have specificity for the various nucleic acid and protein components which it is desired to isolate from a particular sample are described herein and any of these may be used. These distinct solid supports with different surface properties may be provided on different areas of the same solid support, or may be

provided as separate solid supports which are brought into contact with the sample in the same step. These separate solid supports may conveniently be provided for example as separate dipsticks with different appropriate surface properties. Once the desired nucleic acid and protein components have been isolated in such a single step then they can be used and analysed in the same way as if isolated in separate steps.

Once the desired nucleic acid and protein components have been separated onto solid phases these can be subjected to further analysis, e.g. by conventional methods. Again appropriate methods of further analysis will be discussed in more detail below.

The binding of nucleic acid to a solid support is generally independent of its sequence and the specificity of binding is controlled by varying the properties of the solid support and the conditions under which binding of the nucleic acid is induced. As described above the whole nucleic acid component of a sample can be induced to bind to a solid phase or the appropriate choice of binding conditions and/or nature of the solid support can be effected so that DNA or RNA can be selectively bound to a solid phase, thereby enabling a selective DNA or RNA isolation procedure, which may be carried out simultaneously or preferably sequentially. Appropriate methods and conditions whereby all nucleic acid may be bound or DNA and RNA are selectively bound to a solid phase are well known and documented in the art and any of these methods can be used in the methods of the present invention. Some appropriate and preferred methods are discussed briefly here and in more detail below where alternative methods are also discussed.

Conveniently, where DNA and RNA are isolated in two separate steps the sequential separation may take place using two different solid phases, for example solid supports which can differentiate between DNA and RNA.

Thus, such embodiments may comprise carrying out a first separation step to isolate DNA. A further solid support can then be added to the sample to capture the RNA remaining in the sample, either by using a solid support that can bind the RNA, e.g. a silica based support (described in more detail below), or a solid support that can capture specific RNA molecules (e.g. by carrying a complementary nucleic acid probe), or a solid support which can capture a subset of RNA molecules e.g. polyadenylated RNA, using for example a poly-dT or poly-dU based oligonucleotide capture probe. In this way it is possible rapidly to isolate and separate DNA and RNA or subsets of both from the same sample.

In a representative procedure, the sample is lysed in the presence of detergent and the DNA is allowed to bind to a solid support, whereupon the DNA may readily be separated from the sample by removal of the support. Such procedures are described in more detail in WO96/18731 (and in particular Example 12) the disclosure of which is incorporated herein by reference. The procedure of specific DNA capture described in WO96/18731 is also known as the DNA DIRECT method and can be made selective for genomic DNA. Appropriate conditions and methods for carrying out such separations are also described in the Dynabeads DNA DIRECT kit (which is commercially available from Dynal Biotech AS, Oslo, Norway). In methods of the invention where it is desired to isolate DNA and in particular genomic DNA, the DNA DIRECT method of separation is one preferred embodiment. Alternatively, other solid supports having surface chemistries or surface properties supporting DNA binding may be used, for example a support (preferably magnetic beads) carrying surface carboxyl groups.

The RNA may then be isolated from the remaining sample. This can be by a solid phase based system as described above wherein a different solid phase is added to the sample to which RNA becomes specifically bound.

Alternatively the same solid phase may be used and the RNA made to bind thereto by inducing an appropriate change in the sample conditions.

5 Preferably in embodiments where the RNA is isolated by binding to a solid phase, and in the representative procedure as described here, the RNA is mRNA and the surface of the solid phase used has been engineered to carry a capture probe specific for mRNA, e.g. the support has attached to its surface dT oligonucleotides
10 or dU oligonucleotides, e.g. Dynabeads oligo (dT), Dynabeads cDNA release or Dynabeads T₇-oligo(dT) (a new product in which a T7 promoter is included following the principles set out in Ebertine et. al., Biotechniques (1996), 20(4): 584-91 and US-A-5,514,575). The
15 oligonucleotide attached to the "T7 beads" is a combined oligo dT and T7 promoter sequence of the structure 5'AAAAAA-T7sequence(39nt)-dT(20-25)-VN3'.

 Appropriate conditions and methods for carrying out such RNA separations are also described in the product
20 inserts provided with the Dynabeads. In particular Dynabeads oligo (dT) (which are commercially available from Dynal Biotech AS, Oslo, Norway) are sold with an mRNA DIRECT kit protocol which describes how mRNA separation may be carried out. In methods of the
25 invention where it is desired to isolate RNA and in particular mRNA, the mRNA DIRECT method of separation is preferred.

 The choice of oligo dT or dU beads to be used depends on the downstream applications of the isolated
30 mRNA (see further discussion below).

 Support bound oligo dU may be used for isolation of mRNA as discussed in WO 00/58329 of Dynal Biotech AS. This is analogous to the use of oligo dT, but with the added advantage that the "U's" provide a site for
35 cleavage by a glycosylase enzyme.

 In preferred embodiments, the oligo dT/dU (including the T7 modification) constructs additionally

include a further nucleotide sequence (designated "VN", wherein "V" may be any nucleotide other than T (e.g. A, G or C) and "N" may be any nucleotide, including T. This additional "VN" sequence serves the advantageous purpose of assisting in the targeting or positioning of binding of the oligo dU/dT probe sequence to the nucleic acid; the positioning of probe binding is advantageously at the border between the poly A tail and the coding sequence of the mRNA.

10. In the step of the method where protein is isolated, this is conveniently carried out by any appropriate method using a solid phase with appropriate surface properties. Appropriate methods will depend on the type of proteins it is desired to analyse. In one embodiment specific proteins may be isolated using a solid support, which has an appropriate binding partner/ligand attached to its surface, e.g. an antibody to the particular protein which it is desired to isolate. A particularly preferred antibody to be used in this regard is anti-CK19 (e.g. Ex. Dynabeads® anti-CK19 can be used). This thus applies well known principles of affinity separation of proteins, as widely described in the prior art. Any such standard and well known methods may be used or adapted for the present invention.

25 Alternatively a solid phase with more general surface binding properties can be selected, e.g. a solid phase which has surface chemistry which effects classical chromatography interactions such as ion exchange (including both anion exchange and cation exchange), reverse phase interactions or hydrophobic interactions. Such surfaces are conveniently provided on magnetic beads. The use of these more general surfaces conveniently allows the fractionation of proteins in the sample into subsets depending on the structure and properties (charge, hydrophobicity) etc. of the proteins present.

Such general solid phase surfaces can be prepared using conventional techniques which are standard and

well documented in the art of column chromatography. Furthermore, supports (e.g. particles or beads) have been widely described in the art for fluidised bed separation technology, and have properties e.g. ion
5 exchange, designed for protein separation (e.g. US-A-5,084,169, US-A-5,079,155 and US-A-473,231). Magnetic particulate solid supports having such surface chemistry are not known in the art and thus form a further embodiment of the invention. A solid support suitable
10 for protein isolation comprises on its surface positively charged amine groups, which are believed to bind negatively charged proteins. The bound proteins may be eluted under conditions of high salt and low pH. Amine beads are available from Dynal Biotech AS as
15 Dynabeads M-270 amine.

In embodiments of the invention where the protein isolation step is carried out after the various nucleic acid isolation steps, the remaining sample may be subjected to any convenient treatment before the protein
20 is isolated using the appropriate solid support. For example, if it is desired to fractionate the proteins present in the sample, for example using the solid supports with more general surface binding properties as described above, this step can be preceded by a step
25 whereby the protein sample is subdivided into different fractions, e.g. subdivided into one or more of membrane fraction, cytosolic fraction and nuclear fraction. Methods for carrying out such subdivision are standard and well documented in the art of protein analysis.

30 In all the embodiments of the invention described herein, whilst any solid support formats may be used, as well known in the art and widely described in the literature, preferred solid supports are particles and more preferably magnetic particles. Magnetic particles
35 (beads) are well known in the art and widely commercially available. Different sizes of beads are available, or may be prepared, and different

applications (e.g. different DNA, mRNA, protein etc. separations) may be optimised on beads of different sizes, e.g. 4.5 μm , 2.8 μm , 2.7 μm , 1.0 μm in diameter.

5 The methods of the invention are advantageously amenable to automation, particularly if particles, and especially, magnetic particles are used as the support. In a particularly favoured embodiment of the invention, the nucleic acid and protein isolation method is performed using an automated system for handling of the
10 solid support during the cell lysis, nucleic acid binding, protein binding and, optionally, washing steps. Thus the isolated support-bound cells may be transferred to such an apparatus, washed if desired, and lysed; the nucleic acid (i.e. whole nucleic acid
15 component, DNA or RNA) or protein, as appropriate depending on the particular step of the method and the particular support and conditions used may bind to the support, and the bound nucleic acid or protein may readily be washed, using such an apparatus.
20 Furthermore, such an apparatus may also be used to handle the support during the cell/population isolation stage. Thus, it can be seen that the option is there to automate all of the steps of the method if desired.

Particular mention may be made in this regard of
25 the Bead RetrieverTM, available from Dynal ASA, Norway. The apparatus has a system for ready and efficient transfer of the support (carrying cells or nucleic acid or protein) from one well to another. Other automated robotic workstations which may be used include the
30 BioMek 2000 with the Dynal MPC-auto96 magnet station and the TECAN GENESIS RSP with its own built in magnet station. Both the BioMek and the TECAN GENESIS RSP are liquid handling robots.

The present invention provides the first
35 description of a technology which can isolate specific cell populations, DNA, mRNA and proteins from the same sample in an automated way, e.g. by carrying out all the

steps using a solid phase and in particular a magnetic solid phase. Other methods such as laser microdissection are much slower than the methods described herein, are not automatable and require fixed,
5 sectioned material.

The various reactants and components required to perform the methods of the invention may conveniently be supplied in kit form. Such kits represent a further aspect of the invention.

10 At its simplest, this aspect of the invention provides a kit for isolating nucleic acid and protein from the same sample comprising:

(a) a solid support suitable for binding nucleic acid components; and

15 (b) a solid support suitable for binding proteins..

In preferred embodiments the supports of (a) comprise supports which are selective for binding to DNA, or supports which are selective for binding to RNA, or both types of support.

20 Further optional components of the kit include (c) solid supports suitable for isolation of specific cell populations, (d) means for lysing said cells.

The various components (a), (b), (c) and (d) may be as described and discussed above in relation to the
25 methods of the invention.

A further optional component is (e), means for detecting the nucleic acid and/or protein. As discussed above, for detecting nucleic acids such means may include appropriate probe or primer oligonucleotide
30 sequences for use in hybridisation and/or amplification-based detection techniques. For detecting proteins such means include appropriate antibodies.

Optionally further included in such a kit may be buffers, salts, polymers, enzymes etc.

35 Once one or more nucleic acid and protein components have been isolated onto a solid phase, these can then be analysed or used in any appropriate way.

For example, the methods of the invention can be used in any application which involves analysis or comparison of the genome (DNA), transcriptome (RNA) and proteome (expressed protein), or in any analysis where it is
5 desired to compare a particular DNA and/or RNA and the expressed protein, or more preferably a particular DNA, RNA and protein. Such methods are very useful for gene expression studies as they allow analysis and comparison of mRNA levels with protein levels. In addition, the
10 ability to analyse and compare mRNA and protein levels and profiles of samples (e.g. from different cells and tissues) gives a snapshot of the cell's biology.

Furthermore, being able to study the mRNA/protein ratio gives an indication of which genes are subject to post-
15 transcriptional regulation. Such ratios are often shifted during disease states, thereby allowing an indication as to which genes may be involved in disease and allowing a study of the possible regulatory mechanisms behind such a shift in mRNA/protein ratio.

20 The methods of the invention are also useful for example in determining the molecular changes and mechanisms involved in disease states, e.g. determining which genes are mutated, or which proteins are over or under expressed or incorrectly processed. Studying the
25 DNA composition of a sample can provide information on genetic predispositions and acquired mutations, whereas the mRNA and protein profiles generate "molecular portraits" of a state/stage of disease and may also be used for the staging and monitoring of the disease
30 development and treatment. As mentioned above, the methods described are particularly advantageous for such analysis, as they allow the comparison of DNA, RNA and protein from the same sample which is important to allow a more accurate and direct comparison.

35 The method of the invention also has use as a method for isolating nucleic acid and protein fractions from a single sample.

More specifically, in the case of the DNA, once isolated this can be used in nucleic acid based detection procedures, e.g. in genotyping, SNP analysis, sequencing reactions etc.

5 Advantageously the bound nucleic acid need not be eluted or removed from the support prior to carrying out the detection step, although this may be performed if desired. Whether or not the nucleic acid is eluted may also depend on the particular method which was used in
10 the nucleic acid binding step. Thus certain nucleic acid-binding procedures will bind the nucleic acid more tightly than others. In the case of DNA-binding using detergents (e.g. by DNA Direct) for example, the nucleic acid will elute from the solid support when an elution
15 buffer or other appropriate medium is introduced. Nucleic acid bound by means of a precipitant such as alcohol or a chaotrope will remain more tightly bound and may not elute when placed in a buffer medium, and may require heating to be eluted. A preferred use of
20 the DNA is in solid phase SNP analysis, which may optionally be fully automated. Such SNP analysis can be carried out using any appropriate technique. For example, PCR can be carried out using one biotinylated primer, followed by single stranded DNA template
25 generation on streptavidin beads, followed by probe annealing, ddN incorporation and a labelling reaction.

 Thus, the support-bound nucleic acid may be used directly in a nucleic acid based detection procedure, especially if the support is particulate, simply by
30 resuspending the support in, or adding to the support, a medium appropriate for the detection step. Either the nucleic acid may elute into the medium, or as mentioned above, it is not necessary for it to elute.

 A number of different techniques for detecting
35 nucleic acids are known and described in the literature and any of these may be used. Conveniently, nucleic acid may be detected by optical methods, for example by

measuring or determining optical density (OD).
Alternatively, the nucleic acid may be detected by
hybridisation to a probe (which may be a labelled probe
for detection) and very many such hybridisation
5 protocols have been described (see e.g. Sambrook et al.,
1989, Molecular Cloning: A Laboratory Manual, 2nd Ed.
Cold Spring Harbor Press, Cold Spring Harbor, NY). Most
commonly, the detection will involve an *in situ*
hybridisation step, and/or an *in vitro* amplification
10 step using any of the methods described in the
literature for this, for example probe
annealing/hybridization followed by ddN incorporation
(which may be labelled for detection). Thus, techniques
such as LAR, 3SR and the Q-beta-replicase system may be
15 used. However, PCR and its various modifications e.g.
the use of nested primers, or the use of one or more
biotinylated primers will generally be the method of
choice (see e.g. Abramson and Myers, 1993, Current
Opinion in Biotechnology, 4: 41-47 for a review of
20 nucleic acid amplification technologies).

Other detection methods may be based on a
sequencing approach, for example, the minisequencing
approach as described by Syvänen and Söderlund, 1990,
Genomics, 8: 684-692.

25 In amplification techniques such as PCR, the
heating required in the first step to melt the DNA
duplex may release the bound DNA from the support.
Thus, in the case of a subsequent detection step, such
as PCR, the support bound nucleic acid may be added
30 directly to the reaction mix, and the nucleic acid will
elute in the first step of the detection process. The
entire isolated support bound nucleic acid sample
obtained according to the invention may be used in the
detection step, or an aliquot.

35 The results of the PCR or other detection step may
be detected or visualised by many means, which are
described in the art. For example the PCR or other

amplification products may be run on an electrophoresis gel e.g. an ethidium bromide stained agarose gel using known techniques.

5 The amplified nucleic acid may also be detected, or the result confirmed, by sequencing, using any of the many different sequencing technologies which are now available, e.g. standard sequencing, solid phase sequencing, cyclic sequencing, automatic sequencing and minisequencing.

10 If mRNA is isolated, for example using oligo(dT)beads, or oligo(dU) beads (e.g. using Dynabeads oligo(dT), Dynabeads cDNA release or Dynabeads T7 oligo(dT) as described herein) this mRNA can be used or analysed in any appropriate way. The mRNA can for
15 example be analysed using appropriate conventional techniques, e.g. by Northern blotting, or can be used to generate cDNA which may then be subjected to any appropriate DNA based analysis such as those described above.

20 It can be seen that mRNA and cDNA libraries from specific samples, e.g. specific populations or sub-populations of cells, viruses, etc can be produced using the methods of the invention. This application can either be carried out while the mRNA is still on the
25 solid phase, i.e. resulting in solid phase cDNA libraries, or can be carried out after the mRNA has been eluted. The mRNA may also be used as a substrate for RT-PCR. Solid phase cDNA libraries can be used in any appropriate application and can be re-used, for example
30 as template in different PCRs (or other amplifications), thereby enabling PCR (or other) analysis of multiple transcripts, e.g. from single isolated cells. For example, it can be seen from Figure 2 that a solid phase c-DNA library has been re-used in multiple PCR
35 experiments to detect different transcripts in cultured carcinoma cells.

If Dynabeads cDNA release beads (or commercial or

generic equivalents) are used to isolate the mRNA, as described above, these beads have oligo(dU)VN coupled to their surface. These oligo(dU)VN molecules bind to the mRNA and can be used to prime one strand of solid phase cDNA synthesis. The solid phase cDNA can then be released enzymatically by UNG treatment and used, e.g. for probe generation for gene expression profiling.

If Dynabeads T7 oligo(dT) beads (or commercial or generic equivalents) are used to isolate the mRNA, as described above, these beads have coupled to their surface a T7 promoter 5-prime to the oligo(dT)VN. After double stranded solid phase cDNA synthesis (first strand synthesis carried out using the oligo (dT) as a primer and second strand synthesis initiated with random hexamers), the T7 promoter (which is a 39 nucleotide sequence specific for the enzyme T7 polymerase, which recognises this sequence in double stranded DNA) can be used to in vitro transcribe antisense RNA and thereby amplify mRNA from small samples for further analysis, or e.g. for probe generation for gene expression profiling, e.g. for microarray analysis. Such probes are conveniently generated by carrying out reverse transcription of the antisense RNA (with random hexamers) to give cDNA which can be labelled and used to probe microarrays. These cDNAs can also be used in a second round of amplification using T7 beads.

Although microarrays are a powerful high throughput tool, they are not very sensitive - at least 1-2 μ g of mRNA is required per array. This is often impossible to achieve, e.g. by single cell analysis. Thus, in vitro transcription methods are required to amplify the RNA. For this reason the ability to use the Dynabeads T7 oligo(dT) beads to amplify RNA is advantageous in microarray applications and any other applications where amplification of mRNA is advantageous.

Once isolated by binding to a solid phase using the methods of the invention, subsets of proteins (e.g.

those isolated using the supports with more generalised chromatography type surfaces) or individual proteins (e.g. those isolated using supports carrying specific ligands) can be subjected to direct analysis whilst
5 still on the bead (e.g. Bead ELISA) or can be subjected to elution from the solid support followed by analysis by any appropriate conventional technique such as SDS-PAGE/Western blotting, immunoprecipitation, etc.

As discussed extensively above, one of the main
10 advantages of the method of the invention is that one or more of DNA, RNA and protein can be isolated from the same sample ready for analysis. Thus, results obtained by analysing the DNA, RNA and protein from the same sample using any of the individual techniques described
15 above can be compared as appropriate depending on the aim of the analysis.

If the initial sample contains cells or other biological particles which contain the nucleic acid and protein which is to be analysed, then generally the
20 cells/particles to be analysed will need to be separated from the remainder of the sample. Such separation can be carried out by any of the methods which are standard and conventional in the art, such as sedimentation or centrifugation or cell sorting procedures.

Alternatively affinity based separation systems may be used and these are described in more detail below. Such affinity based systems are especially preferred if it is desired to analyse the nucleic acid and protein from a sub-population of the cells or particles contained
30 within the initial sample. Where such affinity-based systems are used, the separation is generally carried out by binding the cells or particles to a solid support carrying an appropriate binding partner.

Affinity-based separation or isolation systems for
35 desired target cells are well known in the art, and rely on the specificity of a binding partner, specific or selective for the target cell, to achieve selective

isolation of the cell. Such a system is employed as an optional step according to the present invention in order to achieve selective isolation of one or more particular populations of cells or other biological "particles" from the sample before it is treated in order to free the nucleic acid and protein and provide the nucleic acid and protein containing sample. Thus, a suitable binding partner for use in this regard may be one or more moieties having a binding affinity for the desired cell or other population(s).

The binding partner may be any molecule or moiety capable of binding to the required populations of cells or other particles, but conveniently will be a protein, polypeptide or peptide. Other moieties or molecules of a different chemical nature, for example carbohydrates or small organic molecules may however also be used. Nucleic acid binding partners e.g. aptamers may also be used.

The binding partner may bind to molecules or structures present on the surface of the cells or other particles, for example to cell surface antigens which are expressed (e.g. specifically) on the surface (e.g. cell specific Dynabeads such as Dynabeads Epithelial Enrich, anti CD14, CD3, CD4, CD8, CD19). Alternatively, the binding partner may be a moiety binding to a cell surface expressed protein e.g. a cell surface receptor.

The binding partner may, for example, conveniently be an antibody specific for a particular surface antigen. Antibody fragments and derivatives may also be used, according to techniques well known in the art. Methods for preparing such fragments or derivatives are well known in the art and widely described in the literature.

Thus, using knowledge of the molecules or other entities present on the surface of the desired population(s), a binding partner, or combination or mixture of binding partners, may be selected to achieve

a desired separation or isolation of cell or other populations from the sample. Advantageously, all or substantially all (i.e. close to all) of the desired cells or other particles present in the sample may be separated. The separation achievable may be dependent not only on the binding partner(s) selected, but also on the nature of the sample, binding conditions etc. Also, biological systems are by their nature variable, and 100% separation may not always be achieved, and, indeed, is not necessary according to the present invention; as in any biological system, some tolerance must be allowed for. However, in preferred embodiments of the invention at least 60%, 65%, 70%, 75%, 80%, 85%, 90% or 95% of the desired populations present in the sample may be separated.

Where more than one different type of binding partner is used they may be attached to the same or different solid supports. Such a system using different solid supports, is applicable particularly in the case of a particulate support such as beads. Thus, different binding partners may be attached to different beads.

In embodiments where more than one different type of binding partner is used, appropriate amounts or ratios at which the different types of binding partner may be used will be readily determined by a person skilled in the art.

As mentioned above, cell separation techniques based on solid phase affinity binding (e.g. immunomagnetic separation (IMS)) are well known in the art and conditions to achieve this may readily be determined by the skilled worker in this field. IMS is a preferred and advantageous cell separation procedure to use, since it enables the isolation of specific cell populations with high purity, for example more than 99% purity from whole blood, as analysed by flow cytometry and molecular techniques. Thus, in the embodiments where a separate affinity isolation step is used, for

example a solid support carrying appropriate binding partner(s) may be brought into contact with the sample. A particulate solid support may, for example, be added to the sample contained (e.g. suspended) in an appropriate medium (e.g. a buffer). The support may then be left in contact with the sample (e.g. incubated) for a length of time to enable binding to the cells or other particles to occur. Conditions during the step are not critical, and the sample-support mixture may be incubated at e.g. 4 to 20°C for 10 minutes to 2 hours e.g. 20-45 minutes. The cell isolation procedure however focuses on minimizing the probability of affecting the physiological state of the cells and thus enabling downstream molecular characterisation and profiling. Laser capture microdissection can also be used to isolate cell populations.

Following cell binding, or any other steps carried out to separate a desired population of cells from which nucleic acid and protein are to be analysed, the isolated or support-bound cells are lysed to release their nucleic acid and protein. Methods of cell lysis are well known in the art and widely described in the literature and any of the known methods may be used. Any of the following methods could, for example, be used: detergent lysis using e.g. SDS, LiDS or sarkosyl in appropriate buffers; the use of chaotropes such as Guanidium hydrochloride (GHC1), Guanidium thiocyanate (GTC), sodium iodide (NaI), perchlorate etc; mechanical disruption, such as by a French press, sonication, grinding with glass beads, alumina or in liquid nitrogen; enzymatic lysis, for example using lysozyme, pronases or cellulases or any of the other lysis enzymes commercially available; lysis of cells by bacteriophage or virus infection; freeze drying; osmotic shock; microwave treatment; temperature treatment; e.g. by heating or boiling, or freezing, e.g. in dry ice or liquid nitrogen, and thawing; alkaline lysis. As

mentioned above, all such methods are standard lysis techniques and are well known in the art, and any such method or combination of methods may be used. The only requirement is that an appropriate method is chosen such that the species of nucleic acid and protein to be analysed in the subsequent procedure remain substantially intact, i.e. substantially non-degraded.

The use of agents such as solvents, alcohols and chaotropes in isolation methods is sometimes disadvantageous. The present invention affords the advantage that the use of such agents may be avoided. Thus, whilst lysis methods such as those mentioned above using such agents may be employed, in advantageous embodiments of the invention the use of such agents is avoided.

Conveniently, lysis may be achieved according to the present invention by using detergents. An exemplary suitable lysis agent thus includes a detergent such as SDS or another alkali metal alkylsulphate salt, e.g. LiDS, or Sarkosyl or combinations thereof. The lysis agents may be supplied in simple aqueous solution, or they may be included in a buffer solution, to form a so-called "lysis buffer". Any suitable buffer may be used, including for example Tris, Bicine, Tricine and phosphate buffers. Alternatively the lysis agents may be added separately. Salts, for example LiCl and NaCl, may also be included in or added to the lysis buffers. In particular, LiCl is preferred when LiDS is used and NaCl is preferred when SDS is used.

Suitable concentrations and amounts of lysis agents will vary according to the precise system etc. and may be appropriately determined, but concentrations of e.g. 2M to 7M chaotropes such as GTC, GHCl, NaI or perchlorate may be used, 0.1M to 1M alkaline agents such as NaOH, and 0.1 to 50% (w/v) e.g. 0.5 to 15% detergent.

To carry out the method of the invention, the support-bound or otherwise isolated cells, may

conveniently be removed or separated from the remainder of the sample, thereby concentrating or enriching the cells. To facilitate subsequent steps, it may be desirable, prior to the lysis step, to dilute the isolated cells, e.g. in an appropriate buffer or other medium. If desired the cells may further be treated, e.g. by heating or mixing (e.g. vortexing). A dilution step may be advantageous to prevent agglomeration/aggregation of a particulate support such as beads. Lysis then may conveniently be achieved by adding an appropriate lysis buffer containing the desired lysis agents or by subjecting the isolated cells to the desired lysis conditions. For example, in the case of simply adding a lysis buffer containing appropriate lysis agents, the isolated cells may simply be incubated in the presence of the lysis buffer for a suitable interval to allow lysis to take place. Different incubation conditions may be appropriate for different lysis systems, and are known in the art. For example for a detergent containing lysis buffer, incubation may take place at room temperature or at higher temperatures e.g. 37°C, 50°C or 65°C. Likewise, time of incubation may be varied from a few minutes e.g. 5 or 10 minutes to hours, e.g. 20 to 40 minutes or 1 to 2 hours. For enzymatic lysis, longer treatment times may be required, e.g. overnight.

Following the obtaining of a sample containing free or released nucleic acid and protein (e.g. following lysis, if required), the steps to be taken will depend on the particular type of analysis it is desired to carry out. For example, in the methods of the invention, solid phases can be used to isolate nucleic acid and protein, DNA and protein, RNA and protein or most preferably DNA, RNA and protein from the same sample. Which entity is isolated depends on the nature of the solid support added and/or the conditions to which the sample and nucleic acid is exposed.

Appropriate conditions and solid supports whereby nucleic acids become bound to the solid support are well known and documented in the art and any of these methods may be used in the embodiments and steps of the

5 invention wherein it is desired that all the released nucleic acids (i.e. DNA and RNA) in a sample are bound to a solid support. Conveniently, the nucleic acid is bound non-specifically to the support i.e. independently of sequence. Thus, for example the released nucleic
10 acid may be precipitated onto the support using any of the known precipitants for nucleic acid, e.g. alcohols, alcohol/salt combinations, polyethylene glycols (PEGs) etc. Precipitation of nucleic acids onto beads in this manner is described for example in WO 91/12079. Thus,
15 salt may be added to the support and released nucleic acid in solution, followed by addition of alcohol which will cause the nucleic acid to precipitate.

Alternatively, the salt and alcohol may be added together, or the salt may be omitted. As described
20 above in relation to the cell binding step, any suitable alcohol or salt may be used, and appropriate amounts or concentrations may readily be determined. However, as mentioned above, it is preferred to avoid the use of solvents, alcohols and similar agents. Thus alternative
25 techniques, avoiding the use of such agents are preferred.

In preferred methods of the invention the DNA and RNA are bound to different solid supports in different steps. Again appropriate conditions and solid supports
30 which allow the selective binding of DNA or RNA are well known in the art and any of these techniques can be used. For example, such preferred techniques whereby DNA and RNA can be separated sequentially from the same sample is described in WO96/18731, the teaching of which
35 is incorporated herein by reference.

For example in this regard, by selecting appropriate "nucleic acid binding" conditions (e.g.

appropriate buffer or lysis/binding medium compositions), it may be selected whether to bind DNA released from the cells, or RNA released from the cells to the solid support. Thus, "binding medium"

5 compositions may be selected favouring DNA binding (or more particularly genomic DNA binding) to the solid support. Such binding medium compositions include those mentioned above, those described in the Examples below, and the compositions of WO96/18731. For example, a
10 representative DNA binding medium may include GuHCl and optionally EDTA.

Briefly, DNA is bound to appropriate non-specific solid supports in the presence of various detergents as described in WO 96/18731 of Dynal AS (the so-called "DNA
15 Direct" procedure). Various detergent-based systems for binding nucleic acids to a solid support are described in this publication and may be used according to the present invention. The solid support is then separated from the rest of the sample, e.g. by the use of a
20 magnetic field.

To bind RNA, appropriate medium compositions or conditions are known in the art, or may readily be determined from RNA isolation procedures known in the art, and may include, for example, the buffers and
25 procedures described in EP-A-0389063 and US-A-5,234,809 of Akzo Nobel NV.

Representative RNA-binding compositions may thus include guanidine thiocyanate (GTC) with EDTA.

For RNA binding, the nature of the solid support
30 may be of importance and in particular a "silica" (i.e. comprising silica itself or being based on silica or a silica derivative) solid surface should be used (see further below)

Advantageously, when the method of the invention is
35 used to isolate DNA, it may be combined with a further step separately to isolate the RNA from the sample. Thus, following the procedure discussed above, and

selecting DNA-binding conditions in the nucleic acid binding step (e.g. a lysis/binding or binding medium favouring DNA binding), DNA released from the support-bound cells may be bound to the support, and removed
5 from the sample. RNA, most notably mRNA, released from the leucocytes, remains in the sample (more precisely in the supernatant). This RNA may readily be isolated from the sample using standard procedures, for example by binding to a capture probe, conveniently immobilised
10 (e.g. by binding to a solid support), consisting of oligo dT.

A further solid support can then be added to the sample under conditions such that the RNA in the sample becomes captured. This can be done by for example using
15 a solid support that can bind the RNA or a solid support that can capture specific RNA molecules (e.g. by carrying a complementary nucleic acid probe), or a subset of RNA molecules, e.g. by carrying a capture probe for polyadenylated RNA, i.e. specific for mRNA.

Conveniently, the nucleic acid or DNA binding step may take place simultaneously or concomitantly with the cell lysis step. This may conveniently be achieved using the detergent-based methods of WO96/18731. Thus,
20 for example, an agent or agents for lysis and nucleic acid binding may conveniently be contained in an appropriate medium (e.g. a buffer or other aqueous solution) and added to the support-bound cells. The cells may then be maintained in contact with the medium e.g. incubated (e.g. as described above) to allow lysis
25 and nucleic acid binding to take place. Such a medium may be referred to as a "lysis/binding" medium. A detergent may function as both lysis agent and to assist in the binding of the nucleic acid to the support.

The detergent may be any detergent, and a vast
35 range are known and described in the literature. Thus, the detergent may be ionic, including anionic and cationic, non-ionic or zwitterionic. The term "ionic

detergent" as used herein includes any detergent which is partly or wholly in ionic form when dissolved in water. Anionic detergents have been shown to work particularly well and are preferred. Suitable anionic
5 detergents include for example sodium dodecyl sulphate (SDS) or other alkali metal alkylsulphate salts or similar detergents, sarkosyl, or combinations thereof.

Conveniently, the detergent may be used in a concentration of 0.2 to 30% (w/v), e.g. 0.5 to 30%,
10 preferably 0.5 to 15%, more preferably 1 to 10%. For anionic detergents concentrations of 1.0 to 5% e.g. 0.5 to 5% have been shown to work well.

The detergent may be supplied in simple aqueous solution, which may be alkaline or acidic, or more
15 preferably in a buffer. Any suitable buffer may be used, including for example Tris, Bicine, Tricine, and phosphate buffers. Conveniently, a source of monovalent cations, e.g. a salt, may be included to enhance nucleic acid capture, although this is not necessary. Suitable
20 salts include chloride salts, e.g. sodium chloride, lithium chloride etc. at concentrations of 0.1 to 1M, e.g. 250 to 500 mM. As mentioned above, other components such as enzymes, may also be included.

Other optional components in the detergent
25 composition include chelating agents e.g. EDTA, EGTA and other polyamino carboxylic acids conveniently at concentrations of 1 to 50 mM etc., reducing agents such as dithiotreitol (DTT) or β -mercaptoethanol, at concentrations of for example 1 to 10 mM.

30 Preferred detergent compositions may for example comprise:

100 mM Tris-HCl pH 7.5

10 mM EDTA

35 2% SDS

or:

100mM TrisCl pH 7.5

10mM EDTA

5% SDS

10mM NaCl

or:

5 100mM TrisCl pH 7.5

500mM LiCl

10mM EDTA

1% LiDS

10 Reference is made to WO96/18731 for further details, exemplary reaction conditions etc.

Alternative nucleic acid binding techniques may also be used in order to achieve the step of binding released nucleic acid to the solid support. For
15 example, one such method may take advantage of the well known principle of nucleic acid binding to a silica surface.

Thus, in such an embodiment, the solid support may comprise or consist of a silica or silica-based or
20 derived material. Many such materials are known and described in the art, and the literature is replete with references describing the isolation of nucleic acids by binding to silica surfaces (see e.g. EP-A-0389063 of AKZO N.V., US-A-5,342,931, US-A-5,503,816 and US-A-
25 5,625,054 of Becton Dickinson, US-A-5,155,018 of Hahnemann University, US-A-6,027,945 of Promega Corp. and US-A-5,945,525 of Toyo Boseki KK).

Ionic binding of the nucleic acid to the support may be achieved by using a solid support having a
30 charged surface, for example a support coated with polyamines.

The support which is used in the method of the invention may also carry functional groups which assist in the specific or non-specific binding of nucleic
35 acids, for example DNA binding proteins e.g. leucine zippers or histones or intercalating dyes (e.g. ethidium bromide or Hoechst 42945) which may be coated onto the

support.

Likewise, the support may be provided with binding partners to assist in the selective capture of nucleic acids. For example, complementary DNA or RNA sequences, or DNA binding proteins may be used. The attachment of such proteins to the solid support may be achieved using techniques well known in the art. Conveniently, such nucleic acid-binding partners may be intermixed on the solid support with the anti-leucocyte binding partners.

The solid supports used for any of the steps of the methods described herein may be any of the well known supports or matrices which are currently widely used or proposed for immobilisation, separation etc. These may take the form of particles, sheets, gels, filters, membranes (e.g. nylon membranes), fibres, capillaries, needles or microtitre strips, tubes, plates or wells, etc, combs, pipette tips, micro arrays, chips, or indeed any solid surface material.

Conveniently the support may be made of glass, silica, latex, plastic or any polymeric material. Generally speaking, for isolation of nucleic acids, the nature of the support is not critical and a variety of surface materials may be used. The surface of the solid support may be hydrophobic or hydrophilic. Preferred are materials presenting a high surface area for binding of the cells, and subsequently, of the nucleic acid. Such supports will generally have an irregular surface and may be for example be porous or particulate e.g. particles, fibres, webs, sinters or sieves. Particulate materials e.g. beads are generally preferred due to their greater binding capacity, particularly polymeric beads/particles.

Conveniently, a particulate solid support used according to the invention will comprise spherical beads. The size of the beads is not critical, but they may for example be of the order of diameter of at least 1 and preferably at least 2 μm , and have a maximum

diameter of preferably not more than 10 and more preferably not more than 6 μm . For example, beads of diameter 2.8 μm , 2.7 μm and 4.5 μm have been shown to work well.

5 Monodisperse particles, that is those which are substantially uniform in size (e.g. size having a diameter standard deviation of less than 5%) have the advantage that they provide very uniform reproducibility of reaction. Monodisperse polymer particles produced by
10 the technique described in US-A-4336173 are especially suitable.

Non-magnetic polymer beads suitable for use in the method of the invention are available from Dynal
15 Particles AS (Lillestrøm, Norway; previously Dyno Particles or Dyno Speciality Polymers) as well as from Qiagen, Amersham Pharmacia Biotech, Serotec, Seradyne, Merck, Nippon Paint, Chemagen, Promega, Prolabo, Polysciences, Agowa, Bangs Laboratories and Dyno
Particles or Dyno Speciality Polymers.

20 However, to aid manipulation and separation, magnetic beads are preferred. The term "magnetic" as used herein means that the support is capable of having a magnetic moment imparted to it when placed in a magnetic field, and thus is displaceable under the
25 action of that field. In other words, a support comprising magnetic particles may readily be removed by magnetic aggregation, which provides a quick, simple and efficient way of separating the particles following any cell, nucleic acid and protein binding steps, and is a
30 far less rigorous method than traditional techniques such as centrifugation which generate shear forces which may disrupt cells or degrade proteins or nucleic acids.

Thus, using the method of the invention, the magnetic particles with cells, nucleic acids, or
35 proteins attached may be removed onto a suitable surface by application of a magnetic field e.g. using a permanent magnet. It is usually sufficient to apply a

magnet to the side of the vessel containing the sample mixture to aggregate the particles to the wall of the vessel and to remove the remainder of the sample for further steps.

5 Especially preferred are superparamagnetic particles for example those described by Sintef in EP-A-106873, as magnetic aggregation and clumping of the particles during reaction can be avoided, thus ensuring uniform cell, nucleic acid and protein extraction. The
10 well-known magnetic particles sold by Dynal Biotech ASA (Oslo, Norway, previously Dynal AS) as DYNABEADS, are particularly suited to use in the present invention. Beads are also automation friendly and allow binding at varying salt concentrations and/or pH. Beads also allow
15 the possibility to manipulate binding and elution conditions of single samples.

Functionalised coated particles for use in the present invention may be prepared by modification of the beads according to US patents 4,336,173, 4,459,378 and
20 4,654,267. Thus, beads, or other supports, may be prepared having different types of functionalised surface, for example positively or negatively charged, hydrophilic or hydrophobic.

In embodiments where an affinity-based separation
25 of cells or other particles is used, the binding partner(s) may be attached to the solid support in any convenient way, before or after binding to the cells/particles, according to techniques well known in the art and described in the literature. Thus the
30 binding partner may be attached directly or indirectly to the solid support.

In a convenient embodiment, the binding partner may be attached to the support, prior to contact with the sample. Such attachment may readily be achieved by
35 methods (e.g. coupling chemistries) well known in the art, and conveniently the binding partner is bound directly to the solid support, for example by coating.

However it may also be attached via spacer or linker moieties. The binding partner may be covalently or reversibly attached according to choice.

Alternatively, as mentioned above, the binding partner may first be brought into contact with the sample, to bind to the cells/particles before being attached to the solid support. In this case, the solid support may conveniently carry or be provided with a binding moiety capable of binding to the particular specific binding partner used. Again, such binding systems are well known in the art. For example, the solid surface may carry a (secondary) antibody capable of binding to the particular binding partner (e.g. a polyclonal anti-species antibody).

In an advantageous embodiment of the invention the lysis step of the method (and indeed the other steps of the method involving the use of a solid support) may include a further step involving the addition of a further or extra amount of solid support (also referred to herein as a "second" solid support) to the reaction mixture.

The use of a second solid support has been found to offer advantages in sample collection for example by improving pellet formation and hence isolation of the first solid support. The improved pellet formation may also reduce non-specific binding of substances or entities in the pellet, or in other words reduces contamination of the pellet. Whilst not wishing to be bound by theory it is believed that when only a first solid support is used the isolated nucleic acid binds to the first solid support as a loose, non-compact mesh, thereby resulting in a relatively loose non-compact pellet. However, where a second support is used and particularly when this second solid support comprises particles which are smaller or larger than the first solid support, the second solid support fills in the gaps (or vice versa) in the loose mesh, thereby making

the pellet tighter and more compact thus reducing the tendency to trap contaminating material.

Further details of the nature and characteristics of appropriate second solid supports for use in the methods of the present invention are described in WO 01/94572, the disclosure of which is incorporated herein by reference.

Briefly, the second solid support may be of comparable size and density to the first solid support. Preferably however, the second solid support is of a smaller size than the first solid support. For example, where the supports are particulate the second solid support comprises particles of a smaller diameter (e.g. approximately half the diameter), than those comprising the first solid support. In especially preferred embodiments the first solid support comprises particles of 4.5 μm diameter (e.g. the M450 beads described herein) whereas the second solid support comprises particles of 2.8 μm diameter (e.g. the M280 and M270 beads described herein). Alternatively, the first support may be smaller than the second support and the dimensions described above may be reversed.

Especially preferably the second solid support may take a more active role in the isolation of the nucleic acid and in such cases the second solid phase is capable of binding to nucleic acid, i.e. has nucleic acid binding properties. Preferably therefore the second solid support may be made of glass, silica, latex, plastic or any polymeric material (i.e. an uncoated surface) capable of binding nucleic acid and such a solid support may optionally be functionalised, for example to aid or improve nucleic acid binding. Particularly preferred in this regard are functionalised solid supports which have a surface charge, preferably a negative surface charge. Most preferred are solid supports coated with carboxylic acid functional groups. Such solid supports are commercially available, for

example the M-270 carboxylic acid beads or M-280 Hydroxyl beads manufactured by Dynal Biotech ASA. Preferably the second solid supports are particulate, e.g. beads, and especially preferably are magnetic.

5 The provision of a further or extra amount of solid support after the cell isolation step results in an improved yield of nucleic acid and also makes the elution of nucleic acid from the solid support easier, particularly where solid supports with a negatively
10 charged surface are used. Whilst not wishing to be bound by theory, as described above it is believed that the addition of an extra amount of a "second" solid support improves the compactness of the bead and nucleic acid pellet and particularly where DNA is able to bind
15 to the second solid support, more effective and complete binding of nucleic acid molecules, rather than the nucleic acid molecules being attached to the beads only at one end or being attached to the beads loosely is achieved.

20 The terms "additional" or "extra" or "further" amount when used herein in connection with the addition of a second solid phase, is used to indicate the addition of any amount (by weight) of second solid phase such that the isolation of nucleic acid is improved, for
25 example the yield of isolated nucleic acid is improved. For example the amount of second solid phase added might be the same amount as the amount of first solid phase used or may be up to approximately 3 to 5 times the amount of first solid phase used. Alternatively, the
30 amount of second solid phase added may be less than the amount of first solid phase providing that the isolation of nucleic acid is improved. Preferably the amount of second solid phase used is 0.5 to 3 times the amount of first solid phase.

35 As the "amount" of solid phase refers to the weight of the solid phase, in the preferred embodiments of the invention where the first and second solid phases are

particulate and the particles making up the second solid phase are smaller (or larger) than the particles making up the first solid phase, the number of particles used for the first and second solid phases will generally be different.

The invention will now be described in more detail in the following non-limiting Example, with reference to the drawings in which:

Figure 1 represents a schematic diagram showing a summary of a model system according to the present invention; leukocytes are isolated by positive selection, binding to beads carrying anti-CD45. DNA is isolated from the captured leukocytes and bound onto DNA-binding beads, from which it may be eluted in purified form, e.g. for subsequent analysis. As an alternative, the isolated leukocyte population may be further fractionated by selection for circulating carcinoma cells (SV₄₀ cells; Ber EP4⁺ cells) using beads (Dynabeads® Epithelial Enrich) carrying a specific binding partner for such cells. This fractionated population may also be subjected to genomic DNA isolation as previously, allowing the general (e.g. leukocyte) and sub-fractionated (e.g. SW₄₀) cell populations to be compared. The isolated cell populations may be subjected to mRNA and protein (specific or general fraction) isolation;

Figure 2 is a photograph of a gel showing the results of PCR experiments carried out as described in Example 1 on cDNA obtained from isolated mRNA (isolated from cultured carcinoma cells), in order to detect different transcripts: Lane 1 - ESX (1st round of PCR); lane 2 - Mucin 1; Lane 3 - CK19 (third round of PCR); Lane 4 - CK19 (first (nested) round of PCR); Lane 4 - ESX (second (nested) round of PCR);

Figure 3 is a photograph of a gel showing the results of PCR and RT-PCR experiments carried out as described in Example 1, showing isolation of genomic DNA

(gDNA) and cDNA encoding CK19: Lane 1 - 100 bp DNA ladder 1 μ g; Lane 2 - cDNA CK19 (mRNA isolation)a; Lane 3 - gDNA CK19 (g-DNA isolation)a; Lane 4 - cDNA CK19 (mRNA isolation)b; Lane 5 - gDNA CK19 (gDNA isolation)b; Lane 6 - Positive control; Lane 7 - Negative control;

Figure 4 shows the results of protein fractionation of a cultured carcinoma cell sample as described in Example 1; Figure 4A shows the results of Bead-ELISA following specific protein isolation using anti-CK19; Figure 4B shows the results of SDS-PAGE comparing the cell lysate against a fractionated cell lysate(fractionated by binding to Dynabeads® M270-amine).

Figure 5a shows the results of an IMS separation of B cells by CD19-IMS and T cells by CD3-IMS, CD8-IMS and CD4-IMS, where mRNA from each cell population is subjected to an RT-PCR for CD19 as described in Example 2. The gels compare the amount of CD19 (and hence B cells) present in each sample to investigate unspecific binding in IMS;

Figure 5b shows the results of an IMS separation of monocytes by CD14, as described in Example 2. This figure shows that RT PCR for CD14 gives a positive result, but RT PCR for CD19 or CD4 result in only low levels being detected and hence demonstrates the low levels of unspecific binding for IMS;

Figure 6 shows the results of an IMS separation of carcinoma cells from whole blood (into which they had been spiked), where RT PCR for CK19 gives a positive result indicating the isolation of the carcinoma cells. RT PCRs for CD4, CD19 and CD14 result in only low levels being detected and this again demonstrates that only low levels of unspecific binding occur for IMS;

Figure 7 is a schematic representation of the DNA isolation protocol in Example 3;

Figure 8 shows that DNA obtained from IMS isolated cells is not degraded;

Figure 9a shows the results for protein isolated using ion exchanger beads from a cell lysate;

Figure 9b shows the results for protein isolated using a cation exchanger;

5 Figure 10a shows PCR results for CK19, CD14, CD4 and CD19 for IMS isolated carcinoma cells, where 2000, 200 or 20 cells were spiked into whole blood. CK19 was detected in samples which only represented 20, 2 or 0.2 cells;

10 Figure 10b shows PCR results for CK19 for IMS isolated carcinoma cells where 1, 5 or 10 cells were spiked into whole blood. CK19 was detected in one of two experiments for one cell.

15 Example 1

Combined isolation of nucleic acid and proteins from single samples

20 Experimental procedures:

Cell pellets containing 2×10^5 cultured carcinoma cells (SW₄₈₀) were lysed in 200 μ l lysis-binding buffer (100 mM Tris-HCl pH 7.5, 500 mM LiCl, 10 mM EDTA pH 8.0, 5 mM DTT, 1% LiDS) containing Dynabeads Epithelial Enrich (200 μ g) and Dynabeads M270 carboxylic Acid (300 μ g),
25 which bind the cells and DNA, respectively. DNA is isolated by incubating the mix on a roller at room temperature for 5 minutes. The bead-DNA complex is separated from the remaining lysate, and washed $3 \times 500 \mu$ l
30 in 1xWashing buffer (10 mM Tris-HCl pH 7.5, 150 mM LiCl, 1.0 mM EDTA), followed by elution in 100 μ l elution buffer (10 mM Tris-HCl pH 8.0, 0.01% Tween 20) at 80°C for 5-10 minutes. The DNA is ready for e.g. PCR (1 μ l is more than enough). Reference is made to Fig. 3 which
35 shows gel analysis of PCR products obtained from *inter alia*, isolated genomic DNA.

To the remaining lysate, 20 μ l oligo(dT) beads (Dynabeads® Oligo (dT) 25 can be used) were added and mRNA annealed for 5 minutes at room temperature on roller. Bead-mRNA complex is separated from the remaining lysate, washed (as standard Dynal procedure ie. as indicated by the manufacturer) and resuspended in 100 μ l ice cold 10 mM Tris-HCl, and ready for RT-PCR. The results are shown in Figs. 2 and 3.

The remaining lysate is diluted (200 μ l lysate + 800 μ l 10 mM Tris-HCl pH 8.0) and added to 1.5 mg Dynabeads M270-amine, which have been pre-washed in 10 mM Tris-HCl pH 8.0. The suspension is incubated on a roller for 1 hour at room temperature. The bead-protein complex is then washed 3x100 μ l in 10 mM Tris-HCl pH 8.0, and eluted by boiling all the beads for 2-5 minutes in 10 μ l SDS-PAGE sample buffer (20 mM Tris-HCl pH 8.0, 1 mM EDTA, 2.5% SDS, 5% b-mercapto-ethanol, 0.01% BPB) and 1 μ l is applied on 10-15% SDS-PAGE Phast gel. The results are shown in Fig. 4 (specifically in Figure 4B).

For specific protein isolation, the lysate was diluted: 5 μ l, 10 μ l and 20 μ l lysate in 100 μ l PBS with 0.1% BSA, prior to bead-DELFIA (ie separation of specific CK19 protein using anti-CK19 beads). The results are shown in Figure 4A.

Isolated LMW in Figure 4B refers to experiments where we dissolved Low Molecular Weight standard (MW protein from Amersham Pharmacia) in lysis/Binding buffer to a conc. of 1 μ g/ μ l and mixed 20 μ l of this with 80 μ l 10 mM Tris-HCl, added 1.5 mg amine beads and incubated and washed, eluted and analysed as described for lysate. This shows that the beads bind proteins.

Example 2

Isolation of cell populations by IMS (immunomagnetic separation)

5

The presence/co-isolation of B cells in different populations of T cells isolated by IMS were investigated. CD19, CD3, CD8 and CD4 IMS isolations were carried out, and mRNA was then extracted from the isolated cell populations and cDNA was synthesised. PCR was then carried out using CD19 primers on a dilution series for each sample, representing from 40 000 to 75 cells. The results for this experiment can be seen in Figure 5a, where the sensitivity of the CD19 RT-PCR is indicated in the first gel. The remaining gels demonstrate the low level of unspecific binding in IMS, since CD19 PCR for the T cell populations demonstrated weak bands and hence only a very low level of unspecific B cell binding had occurred.

20

A similar experiment was carried out, demonstrating a low level of unspecific B cell binding or TM cell binding (<0.2% unspecific) when IMS was used with CD14 to isolate monocytes (Figure 5b).

25

Example 3

Spiking of carcinoma cells into a blood sample and the isolation of the carcinoma cells by IMS

30

1 million carcinoma cells were added to 1 ml of whole blood (i.e. 1 million target cells amongst 4 million non target cells). The carcinoma cells were then separated from the blood cells using IMS with Dynabeads®

35

Epithelial Enrich. As in previous examples, mRNA was isolated from the separated cell population and cDNA was synthesised. PCR was then carried out using CK19

primers, which amplify a transcript not present in blood. The isolation of carcinoma cells could be seen from the results of the CK19 PCR. CD14, CD4 and CD19 PCRs were then carried out to show the specificity of the IMS process for isolating carcinoma cells in this experiment rather than blood cells (Figure 6a).

DNA isolation from the IMS isolated carcinoma cells was carried out using the method as described in Example 1 [IS THIS CORRECT?] (as shown schematically in Figure 7) and as can be seen from Figure 8 there was no degradation of the DNA.

Example 4

Protein isolation with ion and cation exchanger beads

Protein fractionation from the IMS isolated carcinoma cells was carried out using 1 μ m ion exchanger beads. The proteins were bound to the beads in a low salt buffer and were eluted from the beads with a high salt buffer [WE NEED DETAILS]. The ion exchanger Dynabeads gave similar results for protein isolation as Amersham Pharmacia Biotech, ion exchangers (Figure 9a). Protein fractionation was also carried out using cation exchanger beads from spiked cells compared to the cell lysate directly. Figure 9b shows the results from this, where the negative control shows the unspecific background from unspiked blood.

Example 5

Scaling down the techniques to isolate reduced numbers of cells

2000, 200 and 20 carcinoma cells were spiked into 1 ml of whole blood. The cell isolation was carried out as

5

15


25

30


REF ID: A66484

1004.

Spiking of
Sw480 cells
into blood sample



MW 1 2 3 4 5



Re-use of Solid-phase
cDNA library in multiple
PCR experiments,
detecting different
transcripts

Cultured Carcinoma Cells


Fig. 2

Lane	no. of cells	Amplicon	Round of PCR
1	10 000	ESX	first (45 cycles)
2	10 000	Mucin1	second (45 cycles)
3	10 000	CK 19	third (30 cycles)
4	10	CK 19	first (nested)
5	10	ESX	second (nested)

Nucleic acids

PCR and RT-PCR

1 2 3 4 5 6 7






1. 100 bp DNA ladder 1 μ g
2. cDNA CK19(m-RNA isolation)a
- 3.g DNA CK19 (g-DNA isolation)a
4. CDNA CK19 (m-RNA isolation)b
- 5 g DNA CK19 (g-DNA isolation)b
6. Positive control
7. Negative control

Fig. 3

Specific protein isolation

Bead ELISA



Protein fractionation

SDS-PAGE (Phast 10-15%)

1 2 3 4 5 6 7





Fig. 4

Detection of B-cells by CD19 RT-PCR

Cell population


No. cells

1: 40000 6: 1250
2: 20000 7: 625
3: 10000 8: 300
4: 5000 9: 150
5: 2500 10: 75




B-cells
Isolated by
CD19-IMS

<0.3% unspecific




T-cells
Isolated by
CD3-IMS

<<0.2% unspecific



T_C-cells
Isolated by
CD8-IMS

<0.3% unspecific



T_H-cells
Isolated by
CD4-IMS

Figure 5a

Detection of "unspecific" cells in isolated monocyte (CD14) cell population





Figure 5b

Functionality of isolated mRNA



Template in RT-PCR:
mRNA representing
10000 carcinoma cells

Figure 6

Principle of DNA isolation step





Figure 7

Spiked Un-spiked Cells



Integrity of isolated DNA:
No degradation

Cell lysate

LMW std.

Dynabeads

APB-Source 15Q/15S

Anion, pH8 Anion, pH10 Cation, pH8 Cation, pH3.5

Anion, pH8 Anion, pH10 Cation, pH8 Cation, pH3.5




Figure 8

Figure 9a

Different cell no. Neg.ctrl. Cell lysate LMW




Figure 9b




Figure 10a




Figure 10b

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☐ FADED TEXT OR DRAWING
- ☒ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☐ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.